

REPORT DOCUMENTATION PAGE

Form Approved

OMB No. 0704-0188

Reporting burden for this form is estimated to be 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, completing and reviewing the collection of information, and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 1995		3. REPORT TYPE AND DATES COVERED Interim	
4. TITLE AND SUBTITLE A Theoretical Treatment of the Absorption and Emission Properties of Cu(II) Porphyrin			5. FUNDING NUMBERS N00014-90-J-1608 G		
6. AUTHOR(S) Krassimir Stavrev and Michael C. Zerner					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Florida Department of Chemistry Gainesville, FL 32611 USA			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research Chemistry Division Code 1113 Arlington, VA 22217-5000			10. SPONSORING / MONITORING AGENCY REPORT NUMBER Technical Report 30		
11. SUPPLEMENTARY NOTES Published: Chem. Phys. Letters (1994) 233, 179-184.					
12a. DISTRIBUTION / AVAILABILITY STATEMENT This document has been approved for public release: its distribution is unlimited.				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) See attached Abstract.					
14. SUBJECT TERMS					
15. NUMBER OF PAGES 15				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified		18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified		19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	
20. LIMITATION OF ABSTRACT SAR					

19950427 015

DTIC QUALITY INSPECTED 8

OFFICE OF NAVAL RESEARCH

GRANT or CONTRACT N00014-90-J-1608

R&T CODE 4131057- - - 01

Technical Report No. 30

A Theoretical Treatment of the Absorption and Emission Properties of Cu(II) Porphyrin

by

Krassimir K. Stavrev and Michael C. Zerner

Prepared for Publication or Published

in the

Chem. Phys. Letters

University of Florida
Department of Chemistry
Quantum Theory Project
Gainesville, FL 32611-8435

April 18, 1995

Accession For	
NTIS	CRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced <input type="checkbox"/>	
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and / or Special
A-1	

Reproduction in whole or in part is permitted for any purpose of
the United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

A Theoretical Treatment of the Absorption and Emission Properties of Cu(II) Porphyrin

by
Krassimir Stavrev and Michael C. Zerner
Quantum Theory Project
Departments of Chemistry and Physics
University of Florida
Gainesville, FL 32611

Abstract: We examine the electronic spectrum of Cu(II) Porphyrin using the Intermediate Neglect of Differential Overlap (INDO) model that has been specifically parametrized for spectroscopy at the configuration interaction level of theory. The isolated molecule is predicted to have a trip-doublet absorption, corresponding to porphyrin triplets, at about $9,000\text{ cm}^{-1}$ and $13,900\text{ cm}^{-1}$. The Q band is calculated at $15,900\text{ cm}^{-1}$, in good accord with experimental values. We predict a normal emission in this case, as is observed. We find that chelating in the fifth position with a strong ligand such as pyridine leads to a state in energy below the Q band that is mostly $d(z^2) \rightarrow d(x^2-y^2)$ but with appreciable $a_{2u}(\pi) \rightarrow d(x^2-y^2)$ character. We postulate that it is the presence of this state that quenches the emission when Cu(II) porphyrins are complexed with strong fifth or fifth and sixth position ligands.

TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (1)*
Chemistry Division, Code 313
800 North Quincy Street
Arlington, Virginia 22217-5000

Dr. Richard W. Drisko (1)
Naval Civil Engineering
Laboratory
Code L52
Port Hueneme, CA 93043

Defense Technical Information Center (2)
Building 5, Cameron Station
Alexandria, VA 22314

Dr. Harold H. Singerman (1)
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198

Dr. James S. Murday (1)
Chemistry Division, Code 6100
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Eugene C. Fischer (1)
Code 2840
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198

Dr. Robert Green, Director (1)
Chemistry Division, Code 385
Naval Air Weapons Center
Weapons Division
China Lake, CA 93555-6001

Dr. Bernard E. Douda (0)
Crane Division
Naval Surface Warfare Center
Crane, Indiana 47522-5000

Dr. Elek Lindner (1)
Naval Command, Control and
Ocean Surveillance Center
RDT&E Division
San Diego, CA 92152-5000

* Number of copies to forward

ABSTRACT DISTRIBUTION LIST

PROF. DOR BEN-AMOTZ
PURDUE UNIVERSITY
DEPARTMENT OF CHEMISTRY
1393 BROWN BUILDING
WEST LAFAYETTE IN 47907-1393

PROF. MAX BERKOWITZ
DEPARTMENT OF CHEMISTRY
UNIVERSITY OF NORTH CAROLINA
CHAPEL HILL, NORTH CAROLINA 27514

DR. DAVID CHANDLER
DEPARTMENT OF CHEMISTRY
UNIVERSITY OF CALIFORNIA, BERKELEY
BERKELEY CA 94720

PROF. MOSTAFA EL-SAYED
DEPARTMENT OF CHEMISTRY
UNIVERSITY OF CALIFORNIA, LOS ANGELES
LOS ANGELES, CALIFORNIA 90024

PROF. JONATHAN G. HARRIS
DEPARTMENT OF CHEMICAL ENGINEERING
MASSACHUSETTS INSTITUTE OF TECHNOLOGY
CAMBRIDGE, MA 02139

PROF. JURGEN KREUZER
DEPARTMENT OF PHYSICS
DALHOUSIE UNIVERSITY
HALIFAX, NOVA SCOTIA B3H 3J5
CANADA

DR. MARK MARONCELLI
DEPARTMENT OF CHEMISTRY
THE PENNSYLVANIA STATE UNIVERSITY
152 DAVEY LABORATORY
UNIVERSITY PARK, PA 16802

PROF. N. YNGVE OHRN
QUANTUM THEORY PROJECT
UNIVERSITY OF FLORIDA
GAINESVILLE FL 32611

DR. T. PAVLOPOULOUS
NAVAL OCEAN SYSTEMS CENTER
CODE 521
SAN DIEGO, CALIFORNIA 92152-5000

PROF. LAURA PHILIPS
DEPARTMENT OF CHEMISTRY
CORNELL UNIVERSITY
ITHACA, NEW YORK 14853

DR. PETER POLITZER
DEPARTMENT OF CHEMISTRY
UNIVERSITY OF NEW ORLEANS
NEW ORLEANS, LOUISIANA 70148

PROF. GERALDINE RICHMOND
DEPARTMENT OF CHEMISTRY
UNIVERSITY OF OREGON
EUGENE, OREGON 97403

PR JACK SIMONS
DEPARTMENT OF CHEMISTRY
THE UNIVERSITY OF UTAH
SALT LAKE CITY UT 84112
4131050

PROF. DEANNE SNAVELY
DEPARTMENT OF CHEMISTRY
BOWLING GREEN STATE UNIVERSITY
BOWLING GREEN, OHIO 43402

DR. DAVID L. VENEZKY
SURFACE CHEMISTRY BRANCH
CHEMISTRY DIVISION
NAVAL RESEARCH LABORATORY
WASHINGTON DC 20375-5000

PROF. GREGORY A. VOTH
UNIVERSITY OF PENNSYLVANIA
DEPARTMENT OF CHEMISTRY
PHILADELPHIA PA 19104-6323

PROF. GEORGE E. WALRAFEN
CHEMISTRY DEPARTMENT
HOWARD UNIVERSITY
LASER-RESEARCH BUILDING
500 COLLEGE STREET, N.W.
WASHINGTON, D.C. 20059

PROF. S. E. WEBBER
DEPARTMENT OF CHEMISTRY
THE UNIVERSITY OF TEXAS AT AUSTIN
AUSTIN, TEXAS 78712-1167

PROF. MICHAEL ZERNER
DEPARTMENT OF CHEMISTRY
THE UNIVERSITY OF FLORIDA
GAINESVILLE, FL 32611

A Theoretical Treatment of the Absorption and Emission Properties of Cu(II) Porphyrin

by
Krassimir Stavrev and Michael C. Zerner
Quantum Theory Project
Departments of Chemistry and Physics
University of Florida
Gainesville, FL 32611

Abstract: We examine the electronic spectrum of Cu(II) Porphyrin using the Intermediate Neglect of Differential Overlap (INDO) model that has been specifically parametrized for spectroscopy at the configuration interaction level of theory. The isolated molecule is predicted to have a trip-doublet absorption, corresponding to porphyrin triplets, at about $9,000\text{ cm}^{-1}$ and $13,900\text{ cm}^{-1}$. The Q band is calculated at $15,900\text{ cm}^{-1}$, in good accord with experimental values. We predict a normal emission in this case, as is observed. We find that chelating in the fifth position with a strong ligand such as pyridine leads to a state in energy below the Q band that is mostly $d(z^2) \rightarrow d(x^2-y^2)$ but with appreciable $a_{2u}(\pi) \rightarrow d(x^2-y^2)$ character. We postulate that it is the presence of this state that quenches the emission when Cu(II) porphyrins are complexed with strong fifth or fifth and sixth position ligands.

Introduction:

Cu(II) porphyrins (Cu(II)P), figure 1, have been studied extensively both experimentally and theoretically[1-9]. Among the interesting spectroscopic features of these complexes, a luminescence quenching has been observed when chelating in the fifth position with pyridine (*pyr*) and other σ -donor ligands[1-3]. This effect has been attributed to the presence of low-energy states that can be either of the d-d or charge transfer (CT) type[4]. Controversy exists also in the literature regarding the assignment of this mediating state as being of $d-\pi^*$ or $\pi-d$ character. Several models have been proposed to explain the nature of this state[4-9] but none of these has been entirely successful in interpreting the spectroscopy of these complexes.

It is the aim of this work to examine the luminescence quenching observed in these systems through electronic structure calculations on Cu(II)P and Cu(II)P:*pyr* complexes. We use for these studies methods based on SCF/CI as described below, in contrast to methods based on molecular orbital (*mo*) or MS X_α schemes[5-9]. As shown further such an approach is highly successful in reproducing the spectroscopy of Cu porphyrins. It is therefore expected to correctly interpret the origin of the corresponding states of these complexes.

Calculations:

The calculations are performed within the Intermediate Neglect of Differential Overlap model parametrized directly on electronic spectroscopy, INDO/S[10,11]. A self-consistent field (SCF) calculation is performed using the restricted open-shell formalism, ROHF[12]. This calculation is then followed by a configuration interaction calculation (CI) that includes some 1045 singly excited configurations (CIS) for Cu(II)P and some 1126 singly excited configurations for the Cu(II)P-pyridine complex. A valence-bond, or Rumer Diagram, construction is used for this purpose[13,14]. This is essentially all single excitations from orbital 37 to 82 in the former case, where *mo* number 62 is the singly occupied Cu $d(x^2-y^2)$ *mo*, and from orbital 41 to 97, in the latter case, where 77 is the singly occupied *mo*. Three separate calculations have been done on the Cu(II)P-pyridine complex, one in which the Cu to pyridine nitrogen

distance is 2.2 Å, a second in which this distance was set to 2.0 Å and a third in which this distance was set to 1.8 Å. Although the last distance may seem too short to be reasonably obtained, we wanted to examine this extreme asymptote. All calculations were done using the ZINDO program package[15].

Results and Discussion:

The molecular orbital diagram we obtain for the Cu(II)P complex is shown in Figure 2. The labeling is (*mo.* no., symmetry). The mapping of the symmetry labels with these symmetry numbers is given in Table 1. The symmetry labels are given for the subgroup D_{2h} in which the calculation was performed. This is the largest subgroup of the actual symmetry of this complex, D_{4h}. The orbital labeled (62,1) is principally Cu- $d(x^2-y^2)$, is of symmetry b_{1g} in D_{4h} (a_g in D_{2h}), and is singly occupied. The orbital (61,2) is the highest energy occupied, and is of a_{1u}(π) symmetry in D_{4h} (a_u in D_{2h}). The next highest occupied *mo* is (60,4), a_{2u}(π) in D_{4h} (b_{1u} in D_{2h}). The lowest unoccupied *mo*'s are a degenerate pair (63,3) and (64,5), e_g(π^*) in D_{4h} (b_{2g} and b_{3g} in D_{2h}). Note that the singly occupied $d(x^2-y^2)$ *mo* is not the HOMO, as is often the case in SCF calculations involving d localized *mo*'s. As relaxation is very large for such orbitals, nothing should be inferred from the location of this orbital about electron affinities or ionization processes that measure the energy differences between *states*. Removal of an electron from (61,2) and placing it in (62,1) corresponds to a ligand to metal charge transfer excitation and is at considerably higher energy than the state represented in figure 2, see later.

Porphyrin spectra are usually described, at least qualitatively, within the four orbital model of Gouterman[16]. Promotion of an electron from the frontier a_{1u}(π) and a_{2u}(π) orbitals into the e_g(π^*) lead to two close lying E_{1u}($\pi \rightarrow \pi^*$) states with considerable calculated oscillator strength. These two states then interact, leading to a lower lying Q band in which the oscillator strengths tend to cancel, and a higher lying state, the B or Soret, in which the transition moments reinforce. These states are reported in Table 2

as Q and B. The discussion of Q within the four orbital model is reasonably accurate while that of B is only partially complete. Other singly excited states mix with the B band complicating its description, as do higher excitations (double and triple excitations). For these reasons the energy of the Q band is usually calculated in good agreement with experiment, and the B band is calculated some 3,000-4,000 cm^{-1} too high in energy, and with too high an oscillator strength. That is also the case here. The Q band maximum for Cu(II)OEP (octaethyl porphyrin) is reported at 17,670 cm^{-1} in the vapor, to be compared with a calculated value of 15,926 cm^{-1} , while the B band is reported at 25,810 cm^{-1} , 2,700 cm^{-1} lower than our calculated value[17].

As the Cu(II) ion of this complex is a doublet, the overall spin multiplicity of the ground state of the complex is also doublet. The normal singlet manifold of the porphyrin macrocycle is also doublet. The porphyrin triplets also interact with the Cu(II) doublet forming doublets (the trip-doublets of the tables) and quartets (trip-quartets) from the usual rules of the addition of angular momentum[18,19]. The quartet components of these triplet states are calculated at 8,531 cm^{-1} and 13,367 cm^{-1} representing an exchange splitting of 393 cm^{-1} and 530 cm^{-1} respectively.

Ake and Gouterman report emission in Cu(II)TPP (tetraphenyl porphyrin) at 13,300 cm^{-1} which they attribute to the trip-quartet[19], in excellent agreement with our *second* trip-quartet. Although we have found excellent metal-ligand and metal-metal exchange splittings in a variety of systems, CIS calculations of the type performed here using the INDO model have often produced ligand triplet states too low in energy compared to their singlet counterparts, the latter of which are generally in good accord with experiment[20]. It may be that our porphyrin triplets are calculated too low in energy, and thus all states derived from them, trip-doublets and trip-quartets, would also be calculated too low. If it is the *first* trip-quartet that Ake and Gouterman observe, the error in calculated energy for this state would be about 5000 cm^{-1} , a value which seems rather large for the INDO/S model.

In addition to the usual Q and B bands of the porphyrin macrocycle and the trip

doublets that are now spin allowed transitions, there are d->d LaPorte transitions and ligand-to-metal and metal-to-ligand charge transfer transitions. As shown in Table 1 these are all calculated at energies higher than the allowed Q band, and would be difficult to detect as they have no appreciable oscillator strength. The first transitions with predicted oscillator strength are $^2A_g \rightarrow ^2E_u$ calculated at $24,572\text{ cm}^{-1}$, and these would be buried under the B band. The absorption of this complex should be 'normal', while phosphorescence via the trip-doublets might compete with the fluorescence[19].

Cu(II) porphyrin complexes with strong field fifth position ligands do not emit[2]. This by itself suggest the presence of a new state below the Q band that is populated via internal conversion from the Q band, and then to the ground state. Likely candidates for this new state are charge transfer from the Cu(II)P moiety to the fifth position ligand, now tightly bound, or bands that have considerable Cu-d(z^2) orbital participation, as the position of the doubly occupied d(z^2) *mo* should be highly dependent on the presence of fifth and sixth position ligands.

Tables 2 and 3 report the results of calculations on Cu(II)P-pyridine complexes at two different geometries, both with C_{2v} symmetry. At the 2.2 \AA geometry a transition to a state of 2A_1 symmetry that is over 50% d(z^2)->d(x^2-y^2) is calculated just below the Q band. This state is of the same symmetry as is the ground state in C_{2v} , and efficient internal conversion might be expected. At 2.2 \AA this transition has been predicted red shifted by over $6,000\text{ cm}^{-1}$ from its position in the Cu(II)P complex. At the 2.0 \AA geometry, Table 3, this transition is further red shifted by $3,000\text{ cm}^{-1}$. We note that the calculation with Cu(II)-pyridine distance of 1.8 \AA yielded this transition at $7,930\text{ cm}^{-1}$, below both calculated trip-doublets. In all cases this transition is predicted to have considerable ligand-to-metal charge transfer character ($a_{2u}(\pi) \rightarrow d(x^2-y^2)$), a mixing made possible through the reduction in symmetry from D_{4h} to C_{2v} . We attribute the absence of the normal emission to the presence of this state.

Our calculated spectra are presented as figure 3. We note the reproduction in these

calculations of the observed red shift of both Q and B band upon stronger chelation in the fifth position[4]. We also calculate a larger red shift for the B band than for the Q band, also in accord with observed trends[4].

Acknowledgements:

This work was supported in part through grants from the office of Naval Research. We are grateful to the Florida Supercomputer Center at Florida State University for a grant in computer time where most of these calculations were performed. We would like to thank David McMillin (Purdue University) for bringing this problem to our attention.

References:

1. M. Gouterman in "The Porphyrins", D. Dolphin, Ed., Academic Press, New York (1978); Vol.III, Chapter 1.
2. B.E. Smith and M. Gouterman, Chem.Phys. Letters 2 (1968) 517
3. D.R. McMillin, B.P. Hudson, F. Liu, J. Sou, D.J. Bereger and K.A. Meadows, Adv. Chem. Ser. 238 (1993) 211
4. D. Kim, D. Holten and M. Gouterman, J. Am. Chem. Soc. 106 (1984) 2793
5. R.L. Ake and M. Gouterman, Theor. Chim. Acta 15 (1969) 20
6. B. Roos and M. Sundbom, J. Mol. Spectrosc. 36 (1970) 8
7. A. Henriksson, B. Roos and M. Sundbom, Theor. Chim. Acta 27 (1972) 303
8. D. Case and M. Karplus, J. Am. Chem. Soc. 99 (1977) 6182
9. A. Antipas, D. Dolphin, M. Gouterman and E.C. Johnson, J. Am. Chem. Soc. 100 (1978) 7705
10. J.E. Ridley and M.C. Zerner, Theor. Chim. Acta 32 (1973) 111
11. M.C. Zerner, G.H. Loew, R.F. Kirshner and U.T. Mueller-Westerhoff, J. Am. Chem. Soc. 102 (1980) 589
12. W.D. Edwards and M.C. Zerner, Theor. Chim. Acta 72 (1987) 347
13. See, for example, R. Paunz, Spin Eigenfunctions, Plenum Press, New York (1979)
14. J. McKelvey and M.C. Zerner, unpublished
15. ZINDO, a semiempirical program, M.C. Zerner and colleagues, Quantum Theory Project, University of Florida, Gainesville FL 32611
16. M. Gouterman, Chem. Phys. 30 (1959) 1139;
M. Gouterman in "The Porphyrins", D. Dolphin, Ed., Academic Press, New York (1978); Vol.III, Chapter 1.
17. L. Edwards, D. Dolphin and M. Gouterman, J. Mol. Spectrosc. 35 (1970) 90

18. M.G. Cory and M.C. Zerner, Chem. Rev. 91 (1991) 813
19. R. Ake and M. Gouterman, Theor. Chim. Acta 15 (1969) 20; *ibid*
Theor. Chim. Acta 17 (1970) 408
20. J. Ridley and M.C. Zerner, Theor. Chim. Acta 42 (1976) 223

Table 1: Cu(II) Porphyrin reference state 2A_g in D_{2h} . Irreducible representations are: $1=^2A_g$, $2=^2A_u$, $3=^2B_{1g}$, $4=^2B_{1u}$, $5=^2B_{2g}$, $6=^2B_{2u}$, $7=^2B_{3u}$ and $8=^2B_{3g}$.

	ENERGY(CM-1)	(NM)	OSC. STR.	SYM	No. IN IRREP	Comments
1	0.0	0.0	0.0000	1	1	
2	8923.7	1120.6	0.0000	7	2	trip-doub
3	8923.7	1120.6	0.0000	6	2	trip-doub
4	13893.0	719.8	0.0000	7	3	trip-doub
5	13893.1	719.8	0.0000	6	3	trip-doub
6	15926.3	627.9	0.0640	6	4	Q band
7	15926.4	627.9	0.0640	7	4	Q band
8	18390.2	543.8	0.0000	8	2	d(x ² -y ²) - eg(pi*)
9	18390.2	543.8	0.0000	5	2	d(x ² -y ²) - eg(pi*)
10	19489.4	513.1	0.0000	2	2	a _{1u} (pi)-d(x ² -y ²)
11	20281.4	493.1	0.0000	3	2	(pi)- (pi*) mixed
12	21544.8	464.1	0.0000	1	2	d(z ²)-d(x ² -y ²)
13	21668.0	461.5	0.0000	4	2	a _{2u} (pi)-d(x ² -y ²)
14	22261.3	449.2	0.0000	1	3	a _{1u} (pi)- a _{1u} (pi*)
15	22958.1	435.6	0.0000	5	3	d(xz)-eg(pi*)
16	22958.2	435.6	0.0000	8	3	d(yz)-eg(pi*)
17	23269.6	429.7	0.0000	3	3	(pi)-(pi*) mixed
18	24226.3	412.8	0.0000	3	4	(pi)-(pi*) mixed
19	24571.6	407.0	0.0015	6	5	
20	24571.7	407.0	0.0015	7	5	
21	26002.9	384.6	0.0000	3	5	d(xy)-d(x ² -y ²)
22	26194.2	381.8	0.0000	1	4	
23	27504.5	363.6	0.0000	3	6	
24	28566.2	350.1	2.2494	6	6	B band
25	28566.3	350.1	2.2495	7	6	B band

Table 2: Cu(II)Porphyrin + 2.2Å pyridine in C_{2v} symmetry. Irreducible representations are labelled 1 = ²A₁, 2 = ²A₂, 3 = ²B₁ and 4 = ²B₂

	ENERGY(CM-1)	(NM)	OSC. STR.	SYM	No. IN IRREP	COMMENTS
1	0.0	0.0	0.0000	1	1	
2	8477.9	1179.5	0.0000	4	2	trip-doub
3	8531.4	1172.1	0.0000	3	2	trip-doub
4	13007.6	768.8	0.0000	4	3	trip-doub
5	13042.9	766.7	0.0000	3	3	trip-doub
6	15229.7	656.6	0.0000	1	2	50% $d(z^2)$ - $d(x^2-y^2)$
7	15465.2	646.6	0.0445	3	4	Q
8	15530.9	643.9	0.0426	4	4	Q
9	17006.2	588.0	0.0000	4	5	$d(x^2-y^2)$ - eg(π^*)
10	17041.0	586.8	0.0000	3	5	$d(x^2-y^2)$ - eg(π^*)
11	19368.1	516.3	0.0000	2	2	π - π^*
12	19652.9	508.8	0.0000	3	6	$d(xz)$ - $d(x^2-y^2)$
13	20855.3	479.5	0.0000	2	3	$d(xy)$ - $d(x^2-y^2)$
14	21130.3	473.3	0.0001	4	6	$d(yz)$ - $d(x^2-y^2)$
15	21308.7	469.3	0.0000	1	3	pyridine π - π^*
16	21917.8	456.3	0.0000	1	4	π - π^*
17	22107.7	452.3	0.0000	2	4	π - π^*
18	22806.1	438.5	0.0017	4	7	π - π^*
19	22855.8	437.5	0.0018	3	7	π - $d(x^2-y^2)$
20	22973.3	435.3	0.0000	2	5	$a_{1u}(\pi)$ - $d(x^2-y^2)$
21	23273.6	429.7	0.0000	2	6	π - π^*
22	24171.4	413.7	0.0000	1	5	$a_{2u}(\pi)$ - $d(x^2-y^2)$
23	25047.5	399.2	0.0000	1	6	π - π^*
24	26760.6	373.7	0.0104	3	8	$a_{1u}(\pi)$ -pyr(π^*)
25	26800.0	373.1	0.0000	2	7	π - π^*
26	26830.0	372.7	0.0115	3	9	$a_{1u}(\pi)$ -pyr(π^*)
27	27363.1	365.5	1.7434	3	10	B
28	27423.2	364.7	1.6762	4	8	B

Table 3: Cu porphyrin + pyridine at 2.0Å in C_{2v} Symmetry. See table 2 for symmetry labels.

	ENERGY(CM-1)	(NM)	OSC. STR.	SYM	No. IN IRREP	COMMENTS
1	0.0	0.0	0.0000	1	1	
2	8395.7	1191.1	0.0000	4	2	trip-doub
3	8460.3	1182.0	0.0000	3	2	trip-doub
4	12233.5	817.4	0.0000	1	2	a _{2u} (pi)-d(x ² -y ²) + d(z ²)-d(x ² -y ²)
5	12781.7	782.4	0.0000	4	3	trip-doub
6	12802.2	781.1	0.0000	3	3	trip-doub
7	15305.7	653.4	0.0372	3	4	Q
8	15370.3	650.6	0.0340	4	4	Q
9	17092.3	585.1	0.0000	4	5	d(x ² -y ²) -eg(pi*)
10	17120.7	584.1	0.0000	3	5	d(x ² -y ²) -eg(pi*)
11	19223.3	520.2	0.0000	2	2	pi-pi*
12	19491.7	513.0	0.0000	3	6	d(xz)-d(x ² -y ²)
13	20845.9	479.7	0.0000	2	3	d(xy)-d(x ² -y ²)
14	21277.3	470.0	0.0002	4	6	d(yz)-d(x ² -y ²)
15	21287.3	469.8	0.0000	1	3	pyridine pi-pi*
16	21874.0	457.2	0.0000	1	4	pi-pi*
17	21899.2	456.6	0.0000	2	4	pi-pi*
18	22397.2	446.5	0.0030	4	7	pi-pi*
19	22433.5	445.8	0.0032	3	7	pi-pi*
20	22904.9	436.6	0.0000	2	5	a _{1u} (pi)-d(x ² -y ²)
21	22994.7	434.9	0.0000	2	6	pi-pi*
22	23987.9	416.9	0.0000	1	5	a _{2u} (pi)-d(x ² -y ²)
23	24733.6	404.3	0.0000	1	6	pi-pi*
24	26207.1	381.6	0.6429	4	8	split B (35%a _{2u} (pi)->pyr-pi*)
25	26219.6	381.4	0.0850	3	8	
26	26268.1	380.7	0.0893	3	9	
27	26576.5	376.3	1.2947	3	10	B
28	26665.0	375.0	0.0000	2	7	
29	26900.0	371.7	0.7846	4	9	split B (35%a _{2u} (pi)->pyr-pi*)

Figure Captions:

Figure 1: The Cu(II) Porphyrin - pyridine complex of this study, Cu(II)P:pyr .

Figure 2: The frontier molecular orbitals of Cu(II)P. *mo*'s with orbital energies ± 0.05 Hartrees (1 Hartree = 27.21 eV) are plotted as degenerate. The notation (N M) refers to N = *mo* number and M = *mo* symmetry. The assignment of the symmetry numbers to irreducible representations is given in the heading to Table 1. Note that *mo* (62,1) is half-filled. It is not the highest occupied *mo* due to the large amount of relaxation energy associated with doubly occupying this orbital. The excitation (61 2) \rightarrow (62 1) is a charge transfer excitation and lies at higher energy than the configuration designated by this picture, see text.

Figure 3: The simulated spectra of Cu(II)P—, Cu(II)P:pyr at 2.2 Å and Cu(II):pyr at 2.0 Å ---. The vertical lines with no widths shown refer to forbidden excitations. The first taller line at about 8000 cm⁻¹ is the mixed (CT d \rightarrow d) state in Cu(II)P:pyr at 1.8 Å, the shorter four lines mark the location of the trip-doublets, which do not change much upon pyridine chelation. The next taller line is the (CT d \rightarrow d) transition at about 12,500 cm⁻¹ in Cu(II)P:pyr at 2.0 Å, and the line at 15,500 cm⁻¹ the (CT d \rightarrow d) transition in Cu(II)P:pyr at 2.2 Å




Figure 1

0.10		(73,8)	—		$P(\pi^*)$
0.09	(70,4)	(71,1) (72,5)	— — —		$P(\pi^*)$ S $P(\pi^*)$
0.07	(67,5)	(68,8) (69,2)	— — —		$P(\pi^*)$ $P(\pi^*)$ $P(\pi^*)$
0.04		(66,4)	—		$P(\pi^*)$
0.01		(65,2)	—		$P(\pi^*)$
-0.05		(63,5) (64,8)	— —		$P(\pi^*)$ $P(\pi^*)$
-0.24		(61,2)	$\uparrow\downarrow$		$P(\pi)$
-0.26		(60,4)	$\uparrow\downarrow$		$P(\pi)$
-0.33		(58,5) (59,8)	$\uparrow\downarrow$ $\uparrow\downarrow$		$P(\pi)$ $P(\pi)$
-0.34		(57,4) (62,1)	$\uparrow\downarrow$ $\uparrow\downarrow$		$P(\pi)$ Cu d(x ² -y ²)
-0.35	(54,8)	(55,5) (56,4)	$\uparrow\downarrow$ $\uparrow\downarrow$ $\uparrow\downarrow$		$P(\pi)$ $P(\pi)$ $P(\pi)$
-0.41		(53,2)	$\uparrow\downarrow$		$P(\pi)$
	(49,3)	(50,1) (51,7) (52,6)	$\uparrow\downarrow$ $\uparrow\downarrow$ $\uparrow\downarrow$ $\uparrow\downarrow$		Cu d(z ²) $P(\sigma)$ $P(\sigma)$ $P(\sigma)$
-0.46		(48,1)	$\uparrow\downarrow$		$P(\sigma)$
-0.47	(45,5)	(46,8) (47,3)	$\uparrow\downarrow$ $\uparrow\downarrow$ $\uparrow\downarrow$		Cu d(yz) Cu d(xy)
-0.48		(43,7) (44,6)	$\uparrow\downarrow$ $\uparrow\downarrow$		$P(\sigma)$ $P(\sigma)$
-0.52		(41,1) (42,4)	$\uparrow\downarrow$ $\uparrow\downarrow$		$P(\sigma)$ $P(\pi)$

Figure 2

Figure 3

